

Pressure of Hot QCD at Large N_f

Guy D. Moore

*Department of Physics, McGill University,
3600 rue University, Montréal, QC H3A 2T8, Canada*

Abstract

We compute the pressure and entropy of hot QCD in the limit of large number of fermions, $N_f \gg N_c \sim 1$, to next to leading order in N_f . At this order the calculation can be done exactly, up to ambiguities due to the presence of a Landau pole in the theory; the ambiguities are $O(T^8/\Lambda_{\text{Landau}}^4)$ and remain negligible long after the perturbative series (in $g^2 N_f$) has broken down. Our results can be used to test several proposed resummation schemes for the pressure of full QCD.

I. INTRODUCTION

At temperatures large compared to the intrinsic scale of QCD, $T \gg \Lambda_{\text{QCD}}$, the running QCD coupling constant is small, and perturbation theory should be a useful tool for studying thermodynamic and dynamic properties of the theory. This is true in principle, at least.

In practice, there are obstacles to the use of perturbation theory. The first is that hot QCD has an intrinsically nonperturbative scale, at wave numbers $k \sim g^2 T$, or length scales $l \sim 1/g^2 T$. (The running coupling g^2 should be evaluated at a scale of order T .) The presence of nonperturbative physics at this scale has nothing to do with asymptotic freedom or the existence of the scale Λ_{QCD} ; even at temperatures many orders of magnitude above Λ_{QCD} , nonperturbative physics sets in at a scale only a little lower than the thermal scale, $g^2 T \sim T/\log(T/\Lambda_{\text{QCD}})$. Some observables are not very sensitive to this infrared scale; but any observable will be sensitive to this scale at some order in the coupling expansion. For instance, the Debye screening length carries nonperturbative corrections which are suppressed by only one power of g [1, 2]; photon emissivity is believed to receive nonperturbative corrections suppressed by g^2 [3]; and the baryon number violation rate arises entirely from nonperturbative physics, and is sensitive at leading order [4].

The measurables which show thermal effects with the least sensitivity to the nonperturbative scale are the standard thermodynamic parameters, such as pressure, entropy density, and energy density. For these quantities, perturbation theory can determine all coefficients up to $O(g^5)$, and that calculation has been carried out [5, 6, 7]. (It is also possible to compute the $O(g^6 \ln(g))$ term, and this is in progress [8].) With the coefficients of the pressure up to $O(g^5)$ in hand, one can diagnose the quality of the perturbative expansion, by seeing how quickly successive partial sums converge. Here we meet the other problem of perturbation theory at finite temperature; even when the perturbative series exists to some order, it shows very poor convergence. This problem is distinct from the presence of nonperturbative magnetic physics; for instance it also appears in hot QED, which is free of strongly coupled IR behavior [5, 9].

Recently this problem has received renewed attention. Three groups have proposed reorganizations or partial resummations of the thermal perturbative expansion, which are hoped to improve the convergence, and therefore the utility, of perturbation theory. Andersen, Braaten, and Strickland, following an earlier proposal by Karsch *et al.* in scalar field theories [10], have proposed to do so by adding the hard thermal loop (HTL) Lagrangian to the action at tree level and subtracting it again with a counterterm treated as formally higher order, a technique they call hard thermal loop perturbation theory [11, 12, 13]. Blaizot, Iancu, and Rebhan have proposed a technique based on 2PI Φ derivable methods, together with a set of approximations to render the method tractable, also based on hard thermal loops [14, 15, 16]. A similar proposal was recently made by Peshier [17].

All three groups have produced predictions for the QCD pressure, both for pure glue and for QCD with fermions. Unfortunately, it is difficult to judge whether the techniques are successful, since all we know independently about the QCD pressure are the perturbative series, which is only useful at uninterestingly weak coupling, and lattice results from the Bielefeld group [18], which are only at such large coupling that it is not clear whether the resummation techniques should still work. The resummation techniques have also been applied to scalar field theory, but in that case exact results are only known for $O(N)$ scalar

theory at large N , which is an almost trivial theory.

For this reason we think it advantageous to have accurate results for the pressure, in a theory as close to QCD as possible. In this paper we present a calculation of the thermal pressure in QCD with a large number of fermions, $N_f \gg N_c \sim 1$, at next to leading order (NLO) in a large N_f expansion and for general $g^2 N_f$, which is treated as $O(1)$. This theory contains a great deal of the physics of QCD; for instance, the gauge fields are screened by the plasma with the same complicated frequency and momentum dependence as in full QCD. The perturbative series for the pressure shows poor behavior, as in full QCD. Of course, some of the interesting physics of full QCD is absent at NLO in this expansion. This is unfortunate; but on the other hand it is what allows us to solve this theory. For previous exact results in 6 dimensional scalar theory, see [19] (who encountered some similar issues, regarding Landau singularities, to those present here).

We should emphasize now that, since all of the physics of large N_f QCD is also present in full QCD, any approximation which shows poor performance in large N_f QCD cannot be expected to be valid for full QCD either. The reverse need not be the case; an approximation which works at large N_f may stumble over the physics which is left out in the large N_f expansion. Therefore, an approximation's performance against our results should be considered an optimistic estimate of its performance in full QCD.

In the next section, we will explain what we mean by large N_f QCD. In Section III, we describe how the calculation of the QCD pressure is carried out in this theory. Our results and conclusions are presented in Section IV, but we very briefly summarize them here. At small effective coupling $g^2 N_f$, the pressure is close to the free theory value, and perturbation theory works well. As the effective coupling is increased, P/T^4 falls at first, but eventually rises; it becomes of order its free theory value about where Landau pole ambiguities become uncomfortably large. The entropy shows the same behavior. It is difficult to obtain such behavior in an ideal quasiparticle approximation, such as the model of Peshier, Kämpfer, Pavlenko, and Soff [20].

II. LARGE N_f QCD

By large N_f QCD, we mean QCD with the number of colors N_c taken as fixed and $O(1)$, but with the number of fermionic species N_f taken large and the gauge coupling g^2 taken small, so that the combination $g^2 N_f$, which serves as an effective coupling strength, remains fixed and $O(1)$. Therefore, in N_f counting, a factor of g^2 is treated as $O(1/N_f)$. This procedure is appropriate because it is the effective coupling which determines the convergence of perturbation theory.

In the following it will be useful to define g_{eff}^2 as

$$g_{\text{eff}}^2 \equiv \frac{g^2 N_f C_F d_F}{d_A} = \begin{cases} \frac{g^2 N_f}{2}, & \text{QCD,} \\ g^2 N_f, & \text{QED.} \end{cases} \quad (2.1)$$

Here C_F is the quadratic Casimir of the representation containing the fermions and d_F and d_A are the dimensions of the fermionic and adjoint representations. In $SU(N_c)$ gauge

theory but with adjoint fermions, $g_{\text{eff}}^2 = g^2 N_f N_c$. Other than overall factors involving N_c , all the nontrivial coupling dependence of the pressure and entropy which we will find will depend on g_{eff}^2 . Note that g_{eff}^2 is renormalization point dependent; at leading order in N_f its renormalization point dependence is,

$$\beta(g_{\text{eff}}^2) \equiv \frac{\mu dg_{\text{eff}}^2}{d\mu} = \frac{(g_{\text{eff}}^2)^2}{6\pi^2}. \quad (2.2)$$

There are no corrections involving higher powers of g_{eff}^2 ; all corrections to this relation are suppressed by at least one power of $1/N_f$. Therefore, one can solve completely for the scale dependence of g_{eff}^2 , at leading order in $1/N_f$;

$$\frac{1}{g_{\text{eff}}^2(\mu)} = \frac{1}{g_{\text{eff}}^2(\mu')} + \frac{\ln(\mu'/\mu)}{6\pi^2}. \quad (2.3)$$

Clearly the theory has a Landau pole at $\Lambda_{\text{Landau}} \sim \mu e^{6\pi^2/g_{\text{eff}}^2}$. We will define Λ_{Landau} so that the gauge field propagator diverges at $Q^2 = \Lambda_{\text{Landau}}^2$; under this definition, $\Lambda_{\text{Landau}} = \mu e^{5/6} e^{6\pi^2/g_{\text{eff}}^2}$. There is no hope that some strong coupling dynamics somehow generate a strongly coupled UV fixed point, because Eq. (2.3) is exact at leading order in $1/N_f$. Therefore the theory technically does not exist.

This problem is fatal if $\Lambda_{\text{Landau}} \sim T$; but for $\Lambda_{\text{Landau}} \gg T$ we argue that it is not. The Landau pole means that the theory as such becomes ill defined; but it is possible to “rescue” the theory by introducing either additional heavy degrees of freedom or high dimension operators, if they arise at a scale $\leq \Lambda_{\text{Landau}}$. For instance, if the operator $F_{\mu\nu} D^2 D^2 F^{\mu\nu} / \Lambda_{\text{Landau}}^4$ is added to the Lagrangian with coefficient greater than 1, the gauge propagator becomes nonsingular at all energies, which is sufficient for our NLO calculation to proceed. This term ruins the renormalizability of the theory, but in a way which will not appear in the NLO calculation of the pressure. What the Landau pole does mean is, that the exact definition of the theory is ambiguous, and quantities like the entropy or the pressure are only completely well defined after the physics which resolves the Landau singularity has been specified. The size of the ambiguity is, however, suppressed by a positive power of $(T/\Lambda_{\text{Landau}})$, which for the pressure at NLO turns out to be $(T/\Lambda_{\text{Landau}})^4$. So provided that Λ_{Landau} is kept suitably larger than T , the relative size of the ambiguity is tiny and can be ignored.¹ As g_{eff}^2 is increased, this new ambiguity remains tiny long after the convergence of perturbation theory for the pressure has broken down. Therefore, we will forge ahead and compute the pressure, without specifying the UV completion; but we will only work with values of $g_{\text{eff}}^2(\mu)$ for which $(\Lambda_{\text{Landau}}/T) > 35$, which proves abundantly sufficient to keep the relative ambiguity in the NLO pressure below 1%.

The other feature of large N_f gauge theory which we should emphasize is, that the non-abelian nature of the theory is not very important. This is because the gauge field self-interactions carry powers of g without introducing powers of N_f . In particular the gauge self-interactions will have no relevance in the calculation we perform here, and first appear, in the pressure and entropy, at NNLO (next to next to leading order) in $1/N_f$. Therefore our results will apply both to QED and QCD.

¹ Similar considerations, in the context of scalar field theory, appear in Drummond *et al.*, [21].




FIG. 1: Leading order, $O(N_f)$ diagram, (A), which is a free fermion bubble, and NLO, $O(N_f^0)$ diagrams, (B), which are a gauge boson loop with any number of fermion bubble self-energy insertions. (Coupling counterterm insertions, though not shown, should also be resummed.) They can be resummed by the usual Schwinger-Dyson method, to give a loop with a resummed gauge field propagator.

III. COMPUTATION OF THE PRESSURE

In a relativistic field theory with vanishing chemical potentials, the pressure and free energy density are opposites, $P = -F$; so we compute the pressure by computing the free energy, which equals the trace of the sum of all 1PI vacuum bubble graphs. This is to be done at finite temperature, which means in Euclidean space with time extent $\beta = 1/T$, periodic boundary conditions for bosons, and antiperiodic boundary conditions for fermions. To get the *thermal* pressure, we subtract off the result at zero temperature; the difference will be finite, though renormalization is still necessary because the result depends on the coupling g_{eff}^2 and this coupling runs.

At leading order, $O(N_f)$, there is only one diagram, a bare fermionic loop, shown in Fig. 1 (A). Its contribution to the pressure is²

$$P_{\text{LO}} = N_f \text{Tr} \left(\oint_Q^{\text{F}} \ln \mathcal{Q} - \int_Q \ln \mathcal{Q} \right) \quad (3.1)$$

Here the trace is over group and Dirac indices, and the fermionic sum-integral symbol means

$$\oint_Q^{\text{F}} \equiv T \sum_{q^0=(2n+1)\pi T}^{n \in \mathbb{Z}} \int \frac{d^3 \mathbf{q}}{(2\pi)^3}; \quad \int_Q \equiv \int \frac{d^4 Q}{(2\pi)^4}. \quad (3.2)$$

² We use a $[++++]$ Euclidean and $[-+++]$ Minkowski metric. Four-vectors are written with upper case letters, their space and time components in lower case, so if Q is a 4-vector, then q^0 is its time component, $\mathbf{q}$ is the spatial vector and $q \equiv |\mathbf{q}|$ its magnitude. The Minkowski continued frequency is typically denoted with a different letter, usually ω .

We have written this in 3+1 dimensions, but dimensional regularization is implied. At leading order, since the $T = 0$ value is subtracted off, this is irrelevant, the result is finite and equals

$$P_{\text{LO}} = 4N_f d_F \frac{7}{8} \frac{\pi^2 T^4}{90} . \quad (3.3)$$

This is the usual free theory value; $4N_f d_F$ counts the number of degrees of freedom [the 4 counts the two spin states each for particle and anti-particle; d_F is the number of colors, $d_F = N_c$ for fundamental representation fermions in $\text{SU}(N_c)$].

At NLO, there are an infinite number of diagrams, but the structure is extremely simple; all contributions look like a gauge boson propagator with an arbitrary number of fermion loops, and counterterm insertions, inserted. No other graphs contribute at $O(N_f^0)$. This is because the only way to add a loop to a graph without changing the power of N_f , is to add a fermion self-energy loop (or the required counterterm) to a gauge boson propagator; any other way to add a loop introduces a g^2 without another factor of N_f , and $g^2 \sim 1/N_f$. This is the reason that large N_f gauge theory is so simple; the large N_f limit is a much stronger organizing principle than large N_c .

This set of diagrams can of course be resummed by the standard Schwinger-Dyson trick. Therefore the NLO contribution to the pressure is given by

$$P_{\text{NLO}} = -\frac{1}{2} \text{Tr} \left(\oint_Q^{\text{B}} \ln [[G_0^{-1}]^{\mu\nu}(Q) + \Pi_{\text{th}}^{\mu\nu}(Q)] - \int_Q \ln [[G_0^{-1}]^{\mu\nu}(Q) + \Pi_{\text{vac}}^{\mu\nu}(Q)] \right) , \quad (3.4)$$

where the trace runs over group and Lorentz indices. The group trace is trivial, it gives d_A [which is $N_c^2 - 1$ for $\text{SU}(N_c)$ and 1 for QED]. The bosonic sum-integral symbol means

$$\oint_Q^{\text{B}} \equiv T \sum_{Q_0=2n\pi T}^{n \in \mathbb{Z}} \int \frac{d^3 q}{(2\pi)^3} , \quad (3.5)$$

$\Pi_{\text{th}}^{\mu\nu}$ is the fermionic plus counterterm contribution to the gauge boson self-energy at finite temperature, $\Pi_{\text{vac}}^{\mu\nu}$ is the vacuum value, and G_0^{-1} is the free inverse propagator. Our sign convention for Π is chosen so that our Π agrees with the usual Minkowski, $[+ - - -]$ Π . The same expression for the pressure can be derived from the Luttinger-Ward relation [22], with a comparable amount of work.

Both vacuum and thermal self-energies are gauge invariant, since their evaluation doesn't involve gauge field propagators. They are also exactly transverse, $Q_\mu \Pi^{\mu\nu}(Q) = 0$, and together with rotational invariance this ensures that $\Pi_{\text{th}}^{\mu\nu} - \Pi_{\text{vac}}^{\mu\nu}$ can be expressed in terms of two scalar functions [23],

$$\Pi_{\text{th}}^{\mu\nu}(Q) - \Pi_{\text{vac}}^{\mu\nu}(Q) = P^{\mu\nu} \Pi_{\text{T}}(Q) + Q^{\mu\nu} \Pi_{\text{L}}(Q) . \quad (3.6)$$

The exact form of $P^{\mu\nu}$ and $Q^{\mu\nu}$ are not important for our problem, all that matters to us is that they are properly normalized projection operators, $P_\mu^\mu = 2$, $Q_\mu^\mu = 1$, and $P^{\mu\nu}(Q) + Q^{\mu\nu}(Q) + Q^\mu Q^\nu / Q^2 = \eta^{\mu\nu}$ (which is $\delta^{\mu\nu}$, since right now we are in Euclidean space). For the vacuum self-energy, the transverse and longitudinal components are the same, and their value is well known;

$$\Pi_{\text{vac}}^{\mu\nu} = (P^{\mu\nu} + Q^{\mu\nu}) \Pi_{\text{vac}} , \quad \Pi_{\text{vac}} = -\frac{g_{\text{eff}}^2 Q^2}{12\pi^2} \left(\ln \frac{Q^2}{\bar{\mu}^2} - \frac{5}{3} \right) . \quad (3.7)$$




FIG. 2: Points where summation occurs (dots on Euclidean axis) become poles whose residues must be summed. The dashed contour sums these poles, and can be continued (arrows) to the dotted contour, which evaluates the discontinuity across the imaginary (Minkowski) axis.

Here $\bar{\mu}$ is the $\overline{\text{MS}}$ renormalization point.

Some gauge fixing prescription is necessary to evaluate Eq. (3.4), though the result is gauge invariant since it represents a complete treatment to some order in a systematic expansion (this can be explicitly checked). If we choose a gauge where G_0^{-1} factorizes in the same way as Π , the Lorentz trace and logarithm will commute. For instance, in Feynman gauge, $[G_0^{-1}]^{\mu\nu}(Q) = \eta^{\mu\nu}Q^2$, and (dropping the overall group factor)

$$\text{Tr} \ln [[G_0^{-1}]^{\mu\nu}(Q) + \Pi_{\text{th}}^{\mu\nu}(Q)] = 2 \ln(Q^2 + \Pi_T + \Pi_{\text{vac}}) + \ln(Q^2 + \Pi_L + \Pi_{\text{vac}}) + \ln(Q^2) - 2 \ln(Q^2). \quad (3.8)$$

The contribution containing Π_T comes from $P^{\mu\nu}$; the contribution containing Π_L comes from $Q^{\mu\nu}$; the positive $\ln(Q^2)$ contribution comes from $Q^\mu Q^\nu / Q^2$, and the $-2 \ln(Q^2)$ contribution comes from the ghosts, which have no self-energy at this order in N_f . In a general covariant gauge the $Q^\mu Q^\nu / Q^2$ term would be shifted by $\ln(\xi)$ with ξ the gauge fixing parameter; but this contributes equally to the vacuum and thermal contributions and so drops out.³ The next task is to evaluate the thermal self-energy. This has been done by Weldon [23] (in Minkowski space, but the continuation is elementary). His result is listed in the Appendix; unfortunately it involves an integration which to our knowledge cannot be done in closed form.

Because the self-energies Π_L , Π_T are only expressed as single integrals and not as closed form, analytic expressions, at this point we must either make approximations, or turn to semi-numerical methods. We want exact results; otherwise there is no point in our looking at this theory. So the calculation will have to become more numerical. The difference between a sum-integral and an integral is an inconvenient quantity for numerical analysis,

³ This is because $T \sum_{q^0} C = \int (dq^0 / 2\pi) C$ for any C which is q^0 independent. The easiest way to see this is directly in the time domain, though it also follows easily on rotating to Minkowski frequency. The result for P_{NLO} in Coulomb gauge differs by a factor of $\ln(q^2)$ from the covariant result, but this also cancels between vacuum and thermal for the same reason.

especially when issues of UV regulation have not all been addressed. At large values of Q^2 , the argument in the integral in Eq. (3.4) is smooth, and the difference between a sum-integral and an integral is exponentially suppressed, by $O(\exp(-Q/T))$. So it is only the moderate q^2 region where we need to handle sum-integration. In this region, this can be done by re-expressing the frequency sum as a contour integration and continuing to Minkowski frequency. Namely, we note that the function $\cot(q^0/2T)$ has poles at $q^0 = 2n\pi T$, of residue $2T$. So the Matsubara sum can be replaced by a contour integral,

$$\sum_{q^0=2\pi nT}^{n \in \mathbb{Z}} \ln(Q^2 + \Pi(Q)) = i \int_C \frac{dq^0}{4\pi} \cot(q^0/2T) \ln(Q^2 + \Pi), \quad (3.9)$$

where the contour runs just above the real axis from $-\infty$ to $+\infty$, and then just below the axis back to $-\infty$. This is illustrated in Fig. 2, which also shows that this contour can be deformed to pinch the imaginary (Minkowski) axis. The result from upper and lower half planes are equal, so the contour integral becomes

$$\sum_{q^0=2\pi nT}^{n \in \mathbb{Z}} \ln(Q^2 + \Pi(Q)) = - \int_0^\infty \frac{d\omega}{\pi} \coth(\omega/2T) \operatorname{Im} \ln(q^2 - \omega^2 + \Pi(q, i\omega + 0)). \quad (3.10)$$

Note that $\coth(\omega/2T) = 1 + 2/(e^{\omega/T} - 1) = 2(n_b(\omega) + 1/2)$. The $T \rightarrow 0$ limit is trivial; just leave out the Bose distribution and use the vacuum self-energy. Therefore, Eq. (3.4) becomes

$$P_{\text{NLO}} = d_A \int \frac{d^3 q}{(2\pi)^3} \int_0^\infty \frac{d\omega}{\pi} \left[2 \left([n_b + \tfrac{1}{2}] \operatorname{Im} \ln(q^2 - \omega^2 + \Pi_T + \Pi_{\text{vac}}) - \tfrac{1}{2} \operatorname{Im} \ln(q^2 - \omega^2 + \Pi_{\text{vac}}) \right) \right. \\ \left. + \left([n_b + \tfrac{1}{2}] \operatorname{Im} \ln \frac{q^2 - \omega^2 + \Pi_L + \Pi_{\text{vac}}}{q^2 - \omega^2} - \tfrac{1}{2} \operatorname{Im} \ln \frac{q^2 - \omega^2 + \Pi_{\text{vac}}}{q^2 - \omega^2} \right) \right] \quad (3.11)$$

Here Π is always understood to be evaluated at $(q, i\omega + 0)$. Note that $\operatorname{Im} \Pi_{\text{vac}} = (\omega^2 - q^2)(g_{\text{eff}}^2/12\pi)\theta(\omega^2 - q^2)$ is nonzero for timelike momenta; the imaginary parts of the logarithms are always at least $O(g_{\text{eff}}^2)$ away from π . There are no infrared problems in evaluating the above expression; we discuss ultraviolet issues next.

The thermal and vacuum contributions fail to cancel for two reasons. First, there is the factor of $n_b(\omega)$ associated with the thermal contribution, which can be traced to the difference between $\sum_{q^0}$ and $\int dq^0$. There are no ultraviolet problems associated with this piece, since at large ω it is exponentially suppressed by $O(\exp(-\omega/T))$, while at large q but moderate ω , the thermal self-energy has an exponentially small imaginary part, $\operatorname{Im} \Pi \sim \exp(-(q-\omega)/2T)$. Therefore this piece is exponentially ultraviolet safe.

Next, there are Π_T and Π_L . They are only power suppressed at large Q , and can potentially cause more trouble. To understand this region it is best to go back to the Euclidean expression,⁴ and expand in small T^2/Q^2 . In this small T^2/Q^2 limit, the logarithms can be

⁴ We should perform pieces linear in n_b in Minkowski space; but terms without n_b can be performed either in Minkowski space, or in Euclidean space—integrating, not summing, on q^0 —or using some mixture, as convenient.

expanded,

$$2 \ln \frac{Q^2 + \Pi_{\text{vac}} + \Pi_{\text{T}}}{Q^2 + \Pi_{\text{vac}}} + \ln \frac{Q^2 + \Pi_{\text{vac}} + \Pi_{\text{L}}}{Q^2 + \Pi_{\text{vac}}} = \frac{2\Pi_{\text{T}} + \Pi_{\text{L}}}{Q^2 + \Pi_{\text{vac}}} - \frac{2(\Pi_{\text{T}})^2 + \Pi_{\text{L}}^2}{2(Q^2 + \Pi_{\text{vac}})^2} + \dots \quad (3.12)$$

Using Eq. (A1) and Eq. (A7) from the Appendix, the dominant large G term is

$$\frac{1}{2} \int \frac{d^4 Q}{(2\pi)^4} \frac{2\Pi_{\text{T}} + \Pi_{\text{L}}}{Q^2 + \Pi_{\text{vac}}} \simeq \frac{7\pi^2 T^4 g_{\text{eff}}^2}{45} \int \frac{d^4 Q}{(2\pi)^4} \frac{3(q^0)^2 - q^2}{(Q^2)^3} \frac{1}{1 - g_{\text{eff}}^2 \ln[(Q^2)/\bar{\mu}^2]/12\pi^2}. \quad (3.13)$$

This term is potentially logarithmically divergent. If one performs an angular average over the (Euclidean) direction of Q , at fixed $|Q^2|$, then $(3(q^0)^2 - q^2)$ averages to zero, while the rest is a function of Q^2 only; so any regularization or cutoff procedure which respects Euclidean invariance will be UV well behaved. However, if we perform the q^0 integration first, at fixed q , and then perform the $d^3 q$ integration, then because of the logarithm in the denominator, the result does not vanish; instead the q^0 integral gives a result of order $(g_{\text{eff}}^2)^2 T^4 / q^3$. Therefore, performing the q^0 (or Minkowski ω) integration first and the $d^3 q$ integration last holds out the potential for fake logarithmic divergences.

Any sensible way to UV complete the theory, to deal with the Landau pole, will respect Euclidean invariance. Therefore, however we deal with very large Q^2 values, we should do so in a Euclidean invariant way. We choose to apply a large momentum cutoff procedure;⁵ we stop the $d^4 Q$ integration at $Q^2 = a\Lambda_{\text{Landau}}^2$, for $a < 1$, varying the value of a between 1/4 and 1/2 to estimate the irreducible ambiguity. Applying a Euclidean invariant high dimension operator cutoff works similarly.

As we just indicated, it is necessary to cut off the integration at a finite value of Q^2 , lower than $\Lambda_{\text{Landau}}^2$. This also requires some extra care with our Minkowski continuation; we must compute the great arcs at complex q^0 which connect the Minkowski and Euclidean contours. Our procedure is this. First, we perform the Minkowski integration up to $\omega < \Lambda_{\text{Landau}} \sqrt{a}$ and $q < q_{\text{max}} \sim 25T$ (the contribution linear in n_{b} is carried to larger q). The Minkowski integration is done by numerical quadratures, using adaptive mesh refinement. Great care must be taken close to the light-cone at large q . To speed things up, the integrals required for the self-energies are expressed in terms of a handful of integrals of one variable (either $[\omega - q]$ or $[\omega + q]$), which are tabulated and spline interpolated so only two integrations nest. Then we add the great arc from the Minkowski to the Euclidean frequency axis, and integrate over Euclidean q^0 down to $(q^0)^2 + q^2 = q_{\text{max}}^2$. This accounts for all $Q^2 < q_{\text{max}}^2$. Then we integrate over Euclidean $q_{\text{max}}^2 < Q^2 < a\Lambda_{\text{Landau}}^2$, performing first the angular integration and then the integration over the magnitude of Q^2 . Note that $(2\Pi_{\text{T}} + \Pi_{\text{L}})$ in Eq. (3.12) vanishes on angular averaging at least to $O(T^{10})$. The lowest order term at large Q^2 , after angular averaging, comes from $(\Pi_{\text{T}}^2 + \Pi_{\text{L}}^2)/(Q^2 + \Pi_{\text{vac}})^2 \sim T^8$, so the irreducible Landau pole introduced ambiguity is $O(T^8/\Lambda_{\text{Landau}}^4)$, as claimed earlier.

An alternative procedure is to separate the n_{b} and 1/2 pieces of Eq. (3.11) and rotate the 1/2 piece back to Euclidean space. The n_{b} piece is then performed in Minkowski space, where

⁵ The reader may worry that this will introduce gauge fixing dependence. It does not, within the class of Lorentz and Coulomb gauges, presumably because the self-energy is gauge invariant; recall that the nonabelian nature of the theory is not relevant at the order of interest.




FIG. 3: $\mathcal{P} = P_{\text{NLO}}/P_{\text{NLO,free}}$ and $S_{\text{NLO}}/S_{\text{NLO,free}}$ plotted against g_{eff}^2 , left, and $\Lambda_{\text{Landau}}/T$, right. The right plot shows that the features indicating the breakdown of perturbation theory—the pressure bottoming out and then rising—occur long before the temperature reaches Λ_{Landau} .

it is exponentially UV safe; the $1/2$ piece gives Eq. (3.4) but with the frequency summation replaced by a frequency integration; this can directly be done in Euclidean space, rather than by the rather elaborate contour just described.⁶ The answers agree.

We compute the pressure at a series of values of g_{eff}^2 , always at renormalization point

$$\bar{\mu}_{\text{DR}} \equiv \pi e^{-\gamma_E} T \simeq 1.76T, \quad (3.14)$$

with $\gamma_E = 0.577215\dots$ Euler's constant. This is the value suggested by dimensional reduction [24]; at this renormalization point, the energy density in an infrared magnetic field equals its tree level value, $\int d^3x B^2/2$. The result for P_{NLO} is T^4 times a pure function of $g_{\text{eff}}^2(\bar{\mu}_{\text{DR}})$,

$$P_{\text{NLO}} = 2d_A \frac{\pi^2}{90} T^4 \mathcal{P}(g_{\text{eff}}^2(\bar{\mu}_{\text{DR}})). \quad (3.15)$$

We have pulled out the free theory value; $2d_A = 2(N_c^2 - 1)$ counts the number of degrees of freedom in the gauge field, and $\pi^2 T^4/90$ is the usual result for a massless, bosonic field.

One may also compute the entropy density, which is the temperature derivative of the pressure;

$$S_{\text{NLO}} = \frac{dP_{\text{NLO}}}{dT} = 2d_A \frac{2\pi^2}{45} T^3 \left(\mathcal{P} + \frac{(g_{\text{eff}}^2)^2}{24\pi^2} \frac{d\mathcal{P}}{dg_{\text{eff}}^2} \right). \quad (3.16)$$

This can be done from tabulated results for $\mathcal{P}$ by spline interpolation, for instance.

⁶ This simpler procedure was pointed out to me by Ipp and Rebhan.




FIG. 4: Exact result for P_{NLO} , scaled by free value, presented in comparison to the results of perturbation theory. Each figure shows three choices for the renormalization point in perturbation theory; at left, they are within a power of e of $\bar{\mu}_{\text{DR}}$; at right, they are within a factor of 2 of $2\pi T$. The left plot also shows the “best” perturbative result, as explained in the text.

IV. RESULTS AND DISCUSSION

We present our results for the pressure and entropy, plotted against g_{eff}^2 , in Fig. 3. The figure also shows pressure plotted against $\Lambda_{\text{Landau}}/T$. As the coupling is increased, $\mathcal{P}$ at first falls, as expected (for instance in a quasiparticle picture); then it flattens out and rises. The figure shows that the minimum of the pressure occurs when $\Lambda_{\text{Landau}} \simeq 480T$. This is a scale where the Landau singularity is still far off in the ultraviolet; this minimum is a robust prediction of the theory, regardless of the UV completion.

Note that in an earlier (and published!) version of this paper, the numerical results were somewhat different (in particular, higher). This was a result of a coding error in evaluating Eq. (3.11); to take the imaginary part of the log, we used the arctangent of the imaginary over the real piece. For the longitudinal contribution, we forgot to check whether the argument was in the principal range of the arctangent function. Andreas Ipp and Anton Rebhan pointed out a discrepancy between the earlier result and their independent numerical evaluation of the expressions presented here [25], and I thank them for correcting me.

These results are difficult to reconcile with the ideal quasiparticle picture of the plasma presented by Peshier, Kämpfer, Pavlenko, and Soff [20], in which the pressure and entropy are described by treating the plasma as a gas of free but massive quasiparticles. In this model it is unexpected for S_{NLO}/T^3 to rise as a function of g_{eff}^2 , because this requires that the quasiparticle masses get smaller with increasing coupling. Therefore we are doubtful that an ideal quasiparticle picture can be a good description of the physics of large N_f QCD. Since, as we have argued, large N_f is a subset of true QCD, we also don’t expect it to be a

good description of QCD.

We also present the predictions of perturbation theory for the pressure, plotted against the actual value. The perturbative prediction for P_{LO} agrees with Eq. (3.3). The NLO prediction, extracted from the results of Zhai and Kastening [6], is

$$P_{\text{NLO}} = 2d_A \frac{\pi^2 T^4}{90} \left\{ 1 - \frac{25}{2} \frac{g_{\text{eff}}^2}{16\pi^2} + \frac{80}{\sqrt{3}} \left(\frac{g_{\text{eff}}^2}{16\pi^2} \right)^{3/2} + \left[\frac{100}{3} \ln \frac{\mu}{\bar{\mu}_{\text{DR}}} - 68.907 \right] \left(\frac{g_{\text{eff}}^2}{16\pi^2} \right)^2 + \left[\frac{-320}{\sqrt{3}} \ln \frac{\mu}{\bar{\mu}_{\text{DR}}} + \frac{160}{\sqrt{3}} \right] \left(\frac{g_{\text{eff}}^2}{16\pi^2} \right)^{5/2} + O\left((g_{\text{eff}}^2)^3\right) \right\}. \quad (4.1)$$

Note that, unlike in QCD, there is in principle no obstacle to computing the perturbative expansion to any order in g_{eff}^2 . Also there are no appearances of $\ln(g_{\text{eff}}^2)$, because the 3-D dimensionally reduced effective theory is free at order N_f^0 . We have evaluated each partial sum from this series, at a number of renormalization points (running the coupling from $\bar{\mu}_{\text{DR}}$ using the exact relation, Eq. (2.3)), in order to compare the perturbative expansion (and its renormalization point sensitivity) to the exact result. The comparison is shown in Fig. 4. The figure also presents a “best” analytic result. This is what one gets by performing the dimensional reduction step to the highest order known (two loop, in this case) and then solving the dimensionally reduced theory exactly, as advocated recently by Laine [26]. The effective 3 dimensional theory can be solved exactly because it is a free theory. The procedure turns out to be equivalent to evaluating the perturbative series above at the renormalization point where the $(g_{\text{eff}}^2)^{5/2}$ term vanishes. As for QCD, the perturbative expansion is valid at weak coupling but becomes ill behaved long before the temperature reaches the Landau pole (or, in QCD, the phase transition temperature). The “best” analytic result works well to surprisingly large coupling but eventually breaks down.

It is beyond the scope of the current paper to analyze large N_f QCD within each of the proposed resummation schemes, to compare to the exact results presented here. We leave this task to the authors of those schemes. However, to facilitate the comparison, we present our numerical results for the pressure at a number of values of g_{eff}^2 , in Table I.

What are the implications of our results for full QCD? Obviously it does not make sense to use large N_f QCD directly to model the behavior of real world QCD; after all, in the physically interesting range of temperatures, real world QCD has 3 flavors, which is definitely not $\gg N_c = 3$. This was anyway not the point of studying large N_f QCD. Our object was to present a testing ground for other techniques. We expect these techniques to be accurate at small g_{eff}^2 but, possibly, to break down at some larger value. What does that imply for their performance in real QCD? As we have tried to emphasize, success at large N_f probably does not ensure success in full QCD; but failure at large N_f almost certainly means the technique is not valid. But it would also be nice to have a rough way of equating a value of g_{eff}^2 to a comparable value of the QCD coupling α_s , so if a technique works at weak coupling but later breaks down, we can estimate its range of validity in full QCD.

Probably the most reasonable way to equate g_{eff}^2 with α_s is to compare values of the Debye mass. At least, this makes sense in the picture where screening is the dominant physical

g_{eff}^2	$\mathcal{P} = P_{\text{NLO}}/P_{\text{NLO}}(\text{free})$	$\Lambda_{\text{Landau}}/T$
0.0	1.00000	∞
1.0	0.94163	2.1×10^{26}
2.0	0.89799	2.9×10^{13}
3.0	0.86275	1.5×10^9
4.0	0.83364	1.09×10^7
5.0	0.8096	565,000
6.0	0.7897	78,000
7.0	0.7737	19,000
8.0	0.7610	6650
9.0	0.7512	2920
10.0	0.7442	1510
11.0	0.7398	884
12.0	0.7378	564
13.0	0.7381	386
14.0	0.7405	279
15.0	0.7450	210
16.0	0.7515	164
17.0	0.7602	132
18.0	0.7707	109
19.0	0.7832	92
20.0	0.797	78
21.0	0.813	68
22.0	0.832	60
23.0	0.852	53
24.0	0.874	48
25.0	0.898	43

TABLE I: NLO pressure, normalized to the free value, as a function of g_{eff}^2 . The numerical error is less than 2 in the last place shown, and is dominated by numerical issues; Landau pole ambiguity is at most 1 in the last digit in all entries.

process changing the pressure. The Debye mass for each theory, at leading order, is⁷

$$m_{\text{D}}^2(N_{\text{f}} \gg 1) \simeq \frac{g_{\text{eff}}^2}{3} T^2, \quad m_{\text{D}}^2(\text{QCD}) \simeq \frac{2N_{\text{c}} + N_{\text{f}}}{6} g_{\text{s}}^2 T^2 = \frac{2\pi(2N_{\text{c}} + N_{\text{f}})}{3} \alpha_{\text{s}} T^2. \quad (4.2)$$

Equating these, for 3 flavor QCD, gives $g_{\text{eff}}^2 = 18\pi\alpha_{\text{s}}$. So $\alpha_{\text{s}} = 0.11$ (relevant for temperatures of order 100GeV) is comparable to $g_{\text{eff}}^2 = 6$, whereas $\alpha_{\text{s}} = .3$ (relevant closer to the QCD phase transition temperature) is comparable to $g_{\text{eff}}^2 = 17$.

In conclusion, we have presented exact results at next-to-leading order (first nontrivial

⁷ had we chosen $\bar{\mu} = \bar{\mu}_{\text{DR}} e^{1/2}$, the large N_{f} expression would be correct through order $(g_{\text{eff}}^2)^2$.

order) for the pressure of large N_f QCD. These results should be useful for testing the reliability of resummed perturbative techniques, used for full QCD.

ACKNOWLEDGMENTS

I would like to thank Larry Yaffe for presenting this problem to me, and for useful discussions; and Tony Rebhan, for correspondence on the application of the 2PI method to this problem and for reading, and catching an error in, the draft. I also thank Tony Rebhan and Andreas Ipp, for catching a numerical error, which contaminated the original results presented in the first version of this work. This work was supported, in part, by the U.S. Department of Energy under Grant No. DE-FG03-96ER40956 and by a McGill University startup grant.

APPENDIX A: SELF-ENERGIES

The self-energies Π_T , Π_L have been evaluated by Weldon [23]; here we present his results, rotated into Euclidean space, and present a few terms of the expansion about small T^2/Q^2 .

Following Weldon, we define two scalar functions, with normalization chosen to agree with his usage,

$$2g_{\text{eff}}^2 G(Q) \equiv \Pi_\mu^\mu(Q) - [\text{vac}] = 2\Pi_T(Q) + \Pi_L(Q), \quad (\text{A1})$$

$$2g_{\text{eff}}^2 H(Q) \equiv \Pi^{00}(Q) - [\text{vac}] = \frac{q^2}{Q^2} \Pi_L(Q), \quad (\text{A2})$$

so that

$$\Pi_L(Q) = 2g_{\text{eff}}^2 \frac{Q^2}{q^2} H(Q), \quad (\text{A3})$$

$$\Pi_T(Q) = g_{\text{eff}}^2 \left(-\frac{Q^2}{q^2} H(Q) + G(Q) \right). \quad (\text{A4})$$

Weldon's results for these scalar quantities, rotated into Euclidean space and for general Q , are

$$G(Q) = \frac{1}{2\pi^2} \int_0^\infty \frac{dk}{e^{k/T} + 1} \left[4k + \frac{Q^2}{2q} \ln \frac{(q^0)^2 + (2k - q)^2}{(q^0)^2 + (2k + q)^2} \right], \quad (\text{A5})$$

$$H(Q) = \frac{1}{2\pi^2} \int_0^\infty \frac{dk}{e^{k/T} + 1} \left[2k + \frac{Q^2 - 4k^2}{4q} \ln \frac{(q^0)^2 + (2k - q)^2}{(q^0)^2 + (2k + q)^2} + \frac{q^0 k}{q} i \ln \frac{1 + 4k^2/(q^0 - iq)^2}{1 + 4k^2/(q^0 + iq)^2} \right]. \quad (\text{A6})$$

Both expressions are pure real, despite the appearance of i in the expression for $H(Q)$. To obtain the retarded, Minkowski value, replace $q^0 \rightarrow i\omega + 0$.

It is possible to expand each quantity in $T^2 \ll Q^2$ by series expanding the bracketed quantity in small k and then performing the k integration. This is expected to give an

asymptotic series, since the k integration goes up to ∞ but with exponentially decaying weight. The next to leading order result, in Euclidean space, is

$$G(Q \gg T) \rightarrow \frac{7\pi^2(3(q^0)^2 - q^2)}{45(Q^2)^2} T^4 - \frac{248\pi^4(5(q^0)^4 - 10(q^0)^2 q^2 + q^4)}{315(Q^2)^4} T^6 + O(T^8), \quad (\text{A7})$$

$$H(Q \gg T) \rightarrow \frac{7\pi^2 q^2}{45(Q^2)^2} T^4 - \frac{248\pi^4 q^2(5(q^0)^2 - q^2)}{945(Q^2)^4} T^6 + O(T^8). \quad (\text{A8})$$

-
- [1] A. K. Rebhan, Phys. Rev. D **48**, 3967 (1993) [arXiv:hep-ph/9308232].
 - [2] P. Arnold and L. G. Yaffe, Phys. Rev. D **52**, 7208 (1995) [hep-ph/9508280].
 - [3] P. Arnold, G. D. Moore and L. G. Yaffe, JHEP **0111**, 057 (2001) [hep-ph/0109064].
 - [4] See for instance P. Arnold, D. Son and L. G. Yaffe, Phys. Rev. D **55**, 6264 (1997) [hep-ph/9609481]; D. Bödeker, Phys. Lett. B **426**, 351 (1998) [hep-ph/9801430].
 - [5] P. Arnold and C. X. Zhai, Phys. Rev. D **50**, 7603 (1994) [hep-ph/9408276]; Phys. Rev. D **51** (1995) 1906 [hep-ph/9410360].
 - [6] C. X. Zhai and B. Kastening, Phys. Rev. D **52**, 7232 (1995) [hep-ph/9507380].
 - [7] E. Braaten and A. Nieto, Phys. Rev. D **53**, 3421 (1996) [hep-ph/9510408].
 - [8] K. Kajantie, M. Laine, K. Rummukainen, and Y. Schröder, in preparation (private communication).
 - [9] C. Coriano and R. R. Parwani, Phys. Rev. Lett. **73**, 2398 (1994) [arXiv:hep-ph/9405343]. R. R. Parwani, Phys. Lett. B **334**, 420 (1994) [Erratum-ibid. B **342**, 454 (1995)] [arXiv:hep-ph/9406318]. R. R. Parwani and C. Coriano, Nucl. Phys. B **434**, 56 (1995) [arXiv:hep-ph/9409269].
 - [10] F. Karsch, A. Patkos and P. Petreczky, Phys. Lett. B **401**, 69 (1997) [arXiv:hep-ph/9702376].
 - [11] J. O. Andersen, E. Braaten and M. Strickland, Phys. Rev. Lett. **83**, 2139 (1999) [hep-ph/9902327]; Phys. Rev. D **61**, 014017 (2000) [hep-ph/9905337].
 - [12] J. O. Andersen, E. Braaten and M. Strickland, Phys. Rev. D **61**, 074016 (2000) [hep-ph/9908323].
 - [13] J. O. Andersen, E. Braaten, E. Petitgirard and M. Strickland, hep-ph/0205085.
 - [14] J. P. Blaizot, E. Iancu and A. Rebhan, Phys. Rev. Lett. **83**, 2906 (1999) [hep-ph/9906340].
 - [15] J. P. Blaizot, E. Iancu and A. Rebhan, Phys. Rev. D **63**, 065003 (2001) [hep-ph/0005003].
 - [16] J. P. Blaizot, E. Iancu and A. Rebhan, Phys. Lett. B **523**, 143 (2001) [hep-ph/0110369].
 - [17] A. Peshier, Phys. Rev. D **63**, 105004 (2001) [hep-ph/0011250].
 - [18] G. Boyd, J. Engels, F. Karsch, E. Laermann, C. Legeland, M. Lutgemeier and B. Petersson, Nucl. Phys. B **469**, 419 (1996) [hep-lat/9602007]; F. Karsch, E. Laermann and A. Peikert, Phys. Lett. B **478**, 447 (2000) [hep-lat/0002003].
 - [19] D. Bödeker, P. V. Landshoff, O. Nachtmann and A. Rebhan, Nucl. Phys. B **539**, 233 (1999) [arXiv:hep-ph/9806514].
 - [20] A. Peshier, B. Kämpfer, O. P. Pavlenko and G. Soff, Phys. Rev. D **54**, 2399 (1996).
 - [21] I. T. Drummond, R. R. Horgan, P. V. Landshoff and A. Rebhan, Nucl. Phys. B **524**, 579 (1998) [arXiv:hep-ph/9708426].
 - [22] J. M. Luttinger and J. C. Ward, Phys. Rev. **118**, 1417 (1960).

- [23] H. A. Weldon, Phys. Rev. D **26**, 1394 (1982).
- [24] K. Kajantie, M. Laine, K. Rummukainen and M. E. Shaposhnikov, Nucl. Phys. B **458**, 90 (1996) [hep-ph/9508379].
- [25] A. Ipp and A. Rebhan, private communication; A. Ipp, G. D. Moore, and A. Rebhan, hep-ph/0301057, submitted to JHEP.
- [26] M. Laine, arXiv:hep-ph/0301011.